

REFERENCES.

After each reference we mention in brackets the section(s) in which the reference occurs.

- Agrawal, M.K., Coates, J.H., Hunt, D.C. and van der Poorten, A.J. [1980], Elliptic curves of conductor 11, *Math. Comp.* 35, 991–1002. (3.8;3.10; 8.6)
- Alex, L.J. [1976], Diophantine equations related to finite groups, *Comm. Algebra* 4, 77–100. (6.1;6.5)
- Alex, L.J. [1985^a], On the diophantine equation $1 + 2^a = 3^b 5^c + 2^d 3^e 5^f$, *Math. Comp.* 44, 267–278. (1.1;5.4)
- Alex, L.J. [1985^b], On the diophantine equation $1 + 2^a = 3^b 7^c + 2^d 3^e 7^f$, *Arch. Math.* 45, 538–545. (1.1;5.4)
- Babai, L. [1986], On Lovász lattice reduction and the nearest lattice point problem, *Combinatorica* 6, 1–13. (3.4)
- Bachman, G. [1964], *Introduction to p-adic Numbers and Valuation Theory*, Academic Press, New York and London. (2.3)
- Baker, A. [1966], Linear forms in the logarithms of algebraic numbers, *Mathematika* 13, 204–216. (2.4)
- Baker, A. [1968], Contributions to the theory of diophantine equations, I, On the representation of integers by binary forms, II, The diophantine equation $y^2 = x^3 + k$, *Phil. Trans. R. Soc. London*, A 263, 173–208. (8.1)
- Baker, A. [1972], A sharpening of the bounds for linear forms in logarithms I, *Acta Arith.* 21, 117–129. (1.2)
- Baker, A. [1977], The theory of linear forms in logarithms, *Transcendence Theory: Advances and Applications*, Academic Press, London, pp. 1–27. (2.4)
- Baker, A. and Davenport, H. [1969], The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$, *Q. Jl. Math. Oxford* (2) 20, 129–137. (1.1;1.4;3.3)
- Berwick, W.E.H. [1932], Algebraic number fields with two independent units, *Proc. London Math. Soc.* 34, 360–378. (8.2)
- Beukers, F. [1981], On the generalized Ramanujan-Nagell equation, *Acta Arith.* I: 38, 389–410; II: 39, 113–123. (1.1;4.1)

- Billevic^V, K.K. [1956], On the units of algebraic fields of third and fourth degree (Russian), *Mat. Sbornik* Vol. **40**, **82**, 123–137. (8.2)
- Billevic^V, K.K. [1964], A theorem on the units of algebraic fields of n-th degree (Russian), *Mat. Sbornik* Vol. **64**, **106**, 145–152. (8.2)
- Blass, J., Glass, A.M.W., Meronk, D.B. and Steiner, R.P. [1987^a], Practical solutions to Thue equations of degree 4 over the rational integers, *Preprint*, Bowling Green State University. (3.8;8.1)
- Blass, J., Glass, A.M.W., Meronk, D.B. and Steiner, R.P. [1987^b], Practical solutions to Thue equations over the rational integers, *Preprint*, Bowling Green State University. (3.8;8.1;8.4)
- Blass, J., Glass, A.M.W., Meronk, D.B. and Steiner, R.P. [1987^c], Constants for lower bounds for linear forms in the logarithms of algebraic numbers I: The general case, *Preprint*, Bowling Green State University. (2.4)
- Blass, J., Glass, A.M.W., Meronk, D.B. and Steiner, R.P. [1987^d], Constants for lower bounds for linear forms in the logarithms of algebraic numbers II: The homogeneous rational case, *Preprint*, Bowling Green State University. (2.4)
- Borevich, Z.I. and Shafarevich, I.R. [1966], *Number Theory*, Academic Press, New York. (2.1;7.3)
- Bremner, A., Calderbank, R., Hanlon, P., Morton, P. and Wolfskill, J. [1983], Two-weight ternary codes and the equation $y^2 = 4 \times 3^\alpha + 13$, *J. Number Theory* **16**, 212–234. (4.1)
- Brenner, J.L. and Foster, L.L. [1982], Exponential diophantine equations, *Pacific J. Math.* **101**, 263–301. (6.1)
- Brent, R.P. [1978], A Fortran multiple-precision arithmetic package, *ACM Trans. Math. Software* **4**, 57–70; and: Algorithm 524. MP, a Fortran multiple-precision arithmetic package, *ACM Trans. Math. Software* **4**, 71–81. (2.5)
- Brentjes, A.J. [1981], *Multi-dimensional Continued Fraction Algorithms*, MC Tract 145, Centr. Math. Comput. Sci., Amsterdam. (1.3;3.4)
- Brown, E. [1985], Sets in which $xy + k$ is always a square, *Math. Comp.* **45**, 613–620. (1.1)
- Buchmann, J. [1986], A generalization of Voronoi's unit algorithm I & II, *J. Number Theory* **20**, 177–191 & 192–209. (8.2;8.5)
- Buchmann, J. [1987], The generalized Voronoi algorithm in totally real algebraic number fields, to appear. (8.2)
- Cassels, J.W.S. [1957], *An Introduction to Diophantine Approximation*, Cambridge University Press, Cambridge. (1.3)

- Cherubini, J.M. and Wallisser, R.V. [1987], On the computation of all imaginary quadratic fields of class number one, *Math. Comp.* **49**, 295–300. (3.2)
- Coates, J. [1969], An effective p-adic analogue of a theorem of Thue, *Acta Arith.* **15**, 279–305. (2.4;6.1)
- Coates, J. [1970], An effective p-adic analogue of a theorem of Thue II: The greatest prime factor of a binary form, *Acta Arith.* **16**, 399–412. (2.4; 6.1;8.6)
- Delone, B.N. and Faddeev, D.K. [1964], *The theory of irrationalities of the third degree*, Transl. of Math. Monogr., Vol 10, A.M.S., Providence R.I. (8.5)
- Ellison, W.J. [1971^a], Recipes for solving diophantine problems by Baker's method, Sémin. Théorie des Nombres, Université de Bordeaux I, 1970–1, Lab. Th. Nombr. C.N.R.S., Exp. 11, 10 pp. (1.4;3.8)
- Ellison, W.J. [1971^b], On a theorem of S. Sivasankaranarayana Pillai, Sémin. Théorie des Nombres, Université de Bordeaux I, 1970–1, Lab. Th. Nombr. C.N.R.S., Exp. 12, 10 pp. (3.2;5.1)
- Ellison, W.J., Ellison, F., Pesek, J., Stahl, C.E. and Stall, D.S. [1972], The diophantine equation $y^2 + k = x^3$, *J. Number Theory* **4**, 107–117. (3.3;8.1)
- Evertse, J.-H. [1983], *Upper Bounds for the Numbers of Solutions of Diophantine Equations*, MC Tract 168, Centr. Math. Comput. Sci., Amsterdam. (1.1)
- Evertse, J.-H., Györy, K., Stewart, C.L. and Tijdeman, R. [1987], S-unit equations and their applications, *Proceedings of the Durham Symposium on Transcendental Number Theory, July 1986, to appear*. (1.1)
- Faltings, G. [1983], Endlichkeitssätze für abelsche Varietäten über Zahlkörpern, *Invent. Math.* **73**, 349–366. (1.1)
- Fincke, U. and Pohst, M. [1985], Improved methods for calculating vectors of short length in a lattice, including a complexity analysis, *Math. Comp.* **44**, 463–471. (3.6)
- Grinstead, C.M. [1978], On a method of solving a class of diophantine equations, *Math. Comp.* **32**, 936–940. (1.1)
- Hardy, G.H. and Wright, E.M. [1979], *An Introduction to the Theory of Numbers*, (5th ed.), Oxford University Press, Oxford. (1.3;3.2)
- Hasse, H. [1966], Über eine diophantische Gleichung von Ramanujan-Nagell und ihre Verallgemeinerung, *Nagoya Math. J.* **27**, 77–102. (4.1)
- Hunt, D.C. and van der Poorten, A.J., Solving diophantine equations $x^2 + d = a^u$, *unpublished*. (3.2;4.1)

- Kiss, P. [1979], Zero terms in second order linear recurrences, *Math. Sem. Notes Kobe Univ.* (Japan) 7, 145–152. (4.3)
- Knuth, D.E. [1981], *The Art of Computer Programming*, Vol. 2: *Seminumerical Algorithms*, (2nd ed.), Addison-Wesley, Reading Mass. (2.5)
- Koblitz, N. [1977], *p-adic Numbers, p-adic Analysis, and Zeta-functions*, Springer Verlag, New York. (2.3)
- Koblitz, N. [1980], *p-adic Analysis: a Short Course on Recent Work*, Cambridge University Press, Cambridge. (2.3)
- Koksma, J.F. [1937], Diophantische Approximationen, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Vol. 4, Springer Verlag, pp. 407–571. (1.3)
- Lagarias, J.C. and Odlyzko, A.M. [1985], Solving low-density subset sum problems, *J. Assoc. Comput. Mach.* 32, 229–246. (3.4)
- Langevin, M. [1976], Quelques applications de nouveaux résultats de van der Poorten, Séminaire Delange-Pisot-Poitou 1975/76, Paris, Exp. G12, 11 pp. (1.1)
- Lehmer, D.H. [1964], On a problem of Störmer, *Illinois J. Math.* 8, 57–79. (4.9;5.1)
- Lenstra, A.K. [1984], *Polynomial-time Algorithms for the Factorization of Polynomials*, Dissertation, University of Amsterdam. (3.5)
- Lenstra, A.K., Lenstra, H.W. Jr. and Lovász, L. [1982], Factoring polynomials with rational coefficients, *Math. Ann.* 261, 515–534. (1.4; 3.4;3.5)
- Loxton, J.H., Mignotte, M., van der Poorten, A.J. and Waldschmidt, M. [1987], A lower bound for linear forms in the logarithms of algebraic numbers, *C.R. Math. rep. Acad. Sci. Canada* 11, 119–124. (2.4)
- Lutz, É. [1951], *Sur les approximations diophantiennes linéaires P-adiques*, Thèse, Université de Strasbourg. (1.3)
- MacWilliams, F.J. and Sloane, N.J.A. [1977], *The Theory of Error-Correcting Codes*, North-Holland, Amsterdam. (4.1)
- Mahler, K. [1933], Zur Approximation algebraischer Zahlen I: Über den grössten Primteiler binärer Formen, *Math. Ann.* 107, 691–730. (6.1;8.6)
- Mahler, K. [1934], Eine arithmetische Eigenschaft der rekurrenden Reihen, *Mathematica B* (Leiden) 3, 153–156. (4.1)
- Mahler, K. [1935], Über den grössten Primteiler spezieller Polynome zweiten Grades, *Arch. Math. Naturvid. B* 41, 3–26. (4.9;5.1)
- Mahler, K. [1961], *Lectures on Diophantine Approximations I: g-adic Numbers and Roth's Theorem*, University of Notre Dame Press, Notre Dame. (3.10)

- Mignotte, M. [1984^a], On the automatic resolution of certain diophantine equations, *Proceedings of EUROSAM 84*, Lecture Notes in Comput. Sci. 174, Springer Verlag, Berlin, pp. 378–385. (4.1)
- Mignotte, M. [1984^b], Une nouvelle résolution de l'équation $x^2 + 7 = 2^n$, *Rend. Sem. Fac. Sci. Univ. Cagliari* 54, Fasc. 2, 41–43. (4.1)
- Mignotte, M. [1985], $P(x^2+1) \geq 17$ si $x \geq 240$, *C. R. Acad. Sci. Paris* 301, Série I, No. 13, 661–664. (4.9)
- Mignotte, M. and Waldschmidt, M. [1978], Linear forms in two logarithms and Schneider's method, *Math. Ann.* 231, 241–267. (2.4)
- Nagell, T. [1948], Løsning Oppg. 2, 1943, s.29 (Norwegian), *Norsk Mat. Tidsskr.* 30, 62–64. (1.1;4.1;4.9)
- Odlyzko, A.M. and te Riele, H.J.J. [1985], Disproof of the Mertens conjecture, *J. reine angew. Math.* 357, 138–160. (3.4)
- Pethö, A. [1983], Full cubes in the Fibonacci sequence, *Publ. Math. Debrecen* 30, 117–127. (1.1;8.1).
- Pethö, A. [1985], On the solution of the diophantine equation $G_n = p^z$, *Proceedings EUROCAL '85*, Linz, Austria, Vol 2, Lecture Notes in Comput. Sci. 204, Springer Verlag, Berlin, pp. 503–512. (4.1;4.7)
- Pethö, A. and Schulenberg, R. [1987], Effektives Lösen von Thue Gleichungen, *Publ. Math. Debrecen*, to appear. (3.8;8.1)
- Pethö, A. and de Weger, B.M.M. [1986], Products of prime powers in binary recurrence sequences I: The hyperbolic case, with an application to the generalized Ramanujan-Nagell equation, *Math. Comp.* 47, 713–727. (1.1; 2.2;4)
- Philippon, P. and Waldschmidt, M. [1987], Lower bounds for linear forms in logarithms, preprint. (2.4)
- Pinch, R. [1987], Simultaneous Pellian equations, preprint, University of Cambridge. (1.1)
- Pohst, M. and Zassenhaus, H. [1982], On effective computation of fundamental units I & II, *Math. Comp.* 38, 275–291 & 293–329. (8.2)
- van der Poorten, A.J. [1977], Linear forms in logarithms in the p-adic case, *Transcendence Theory: Advances and Applications*, Academic Press, London, pp. 29–57. (1.2;2.4;6.2)
- Rumsey, H. Jr. and Posner, E.C. [1964], On a class of exponential equations, *Proc. Am. Math. Soc.* 15, 974–978. (4.9;6.1)
- Schinzel, A. [1967], On two theorems of Gelfond and some of their applications, *Acta Arith.* 13, 177–236. (2.4;4.1)

- Shorey, T.N., van der Poorten, A.J., Tijdeman, R. and Schinzel, A. [1977],
 Applications of the Gel'fond-Baker method to diophantine equations,
 Transcendence Theory: Advances and Applications, Academic Press, London,
 pp. 59-77. (1.1)
- Shorey, T.N. and Tijdeman, R. [1986], *Exponential Diophantine Equations*,
 Cambridge University Press, Cambridge. (1.1;1.2;4.2;5.1;6.1;8.1;8.6)
- Sprindzuk, V.G. [1969], Effective estimates in 'ternary' exponential
 diophantine equations (Russian), *Dokl. Akad. Nauk BSSR* 12, 293-297.
 (6.1)
- Steiner, R.P. [1977], A theorem on the Syracuse problem, *Proc. Seventh
 Manitoba Conf. Numer. Math. Comp.*, pp. 553-559. (3.2)
- Steiner, R.P. [1986], On Mordell's equation $y^2 - k = x^3$: a problem of
 Stolarsky, *Math Comp.* 46, 703-714. (3.3;8.1)
- Stewart, C.L. and Tijdeman, R. [1986], On the Oesterlé-Masser conjecture,
 Monatsh. Math. 102, 251-257. (6.6)
- Størmer, C. [1897], Quelques théorèmes sur l'équation de Pell $x^2 - Dy^2 = \pm 1$
 et leurs applications, *Vid. Skr. I Math. Natur. Kl. (Christiana)*, 1897,
 No. 2, 48 pp. (4.9;5.1)
- Stroeker, R.J. and Tijdeman, R. [1982], Diophantine equations, (with an
 Appendix by P.L. Cijssouw, A. Korlaar and R. Tijdeman), *Computational
 Methods in Number Theory*, MC Tract 155, Centr. Math. Comp. Sci.,
 Amsterdam, pp. 321-369. (1.1;3.2;5.1;5.4)
- Thue, A. [1909], Über Annäherungswerten algebraischer Zahlen, *J. reine
 angew. Math.* 135, 284-305. (8.1)
- Tijdeman, R. [1973], On integers with many small prime factors, *Compositio
 Math.* 26, 319-330. (5.1)
- Tijdeman, R. [1976], On the equation of Catalan, *Acta Arith.* 29, 197-209.
 (1.1)
- Tijdeman, R. [1985], On the Fermat-Catalan equation, *Jahresber. Deutsche
 Math. Verein.* 87, 1-18. (1.1)
- Tijdeman, R. and Wang, L. [1987], Sums of products of powers of given prime
 numbers, *Preprint*, University of Leiden. (1.1;5.4)
- Tzanakis, N. [1983], On the diophantine equation $y^2 - D = 2^k$, *J. Number
 Theory* 17, 144-164. (4.1)
- Tzanakis, N. [1984], The complete solution in integers of $x^3 + 2y^3 = 2^n$,
 J. Number Theory 19, 203-208. (8.6)
- Tzanakis, N. [1987], On the practical solution of the Thue equation, an
 outline, *Proceedings Colloquium on Number Theory, Budapest, July 1987*,
 to appear. (8)

- Tzanakis, N. and de Weger, B.M.M. [1987], On the practical solution of the Thue equation, *Memorandum No. 668*, Faculty of Applied Mathematics, University of Twente, to appear. (1.1;8)
- Tzanakis, N. and Wolfskill, J. [1986], On the diophantine equation $y^2 = 4q^n + 4q + 1$, *J. Number Theory* 23, 219–237. (4.1)
- Tzanakis, N. and Wolfskill, J. [1987], The diophantine equation $x^2 - 4q^{a/2} + 4q + 1$ with an application in coding theory, to appear. (4.1)
- Vojta, P. [1987], *Diophantine Approximations and Value Distribution Theory*, Lecture Notes in Mathematics 1239, Springer Verlag, Berlin. (6.6)
- Wagstaff, S.S. Jr. [1979], Solution of Nathanson's exponential congruence, *Math. Comp.* 33, 1097–1100. (3.9)
- Wagstaff, S.S. Jr. [1981], The computational complexity of solving exponential congruences, *Congressus Numerantium*, University of Winnipeg, Canada, Vol. 31, pp. 275–286. (3.9)
- Waldschmidt, M. [1980], A lower bound for linear forms in logarithms, *Acta Arith.* 37, 257–283. (2.4)
- de Weger, B.M.M. [1986^a], Approximation lattices of p-adic numbers, *J. Number Theory* 24, 70–88. (1.4;3.10)
- de Weger, B.M.M. [1986^b], Products of prime powers in binary recurrence sequences II: The elliptic case, with an application to a mixed quadratic-exponential equation, *Math. Comp.* 47, 729–739. (1.1;4)
- de Weger, B.M.M. [1987^a], Solving exponential diophantine equations using lattice basis reduction algorithms, *J. Number Theory* 26, 325–367. (1.1;5;6)
- de Weger, B.M.M. [1987^b], On the practical solution of Thue–Mahler equations, an outline, *Memorandum 649*, Faculty of Applied Mathematics, University of Twente. (1.1;8.6)
- Yu, K.R. [1987^a], Linear forms in the p-adic logarithms, *Report MPI/87-20*, Max Planck Institut für Mathematik, Bonn. (1.2;2.4;6.2)
- Yu, K.R. [1987^b], Linear forms in logarithms in the p-adic case, *Proceedings of the Durham Symposium on Transcendental Number Theory*, July 1986, to appear. (2.4)